

CoLyte Optical Colonoscopy Instruction

PROCEDURE DATE

ARRIVAL TIME (*Not Procedure Time*)

PURPOSE OF COLONOSCOPY:

- (A) Abnormalities can be visualized and further information obtained by taking a biopsy.
- (B) A diagnostic exam that allows the physician to carefully examine the lining of the large intestine or colon with a flexible tube equipped with a light and a camera.
- (C) A therapeutic exam in that if a polyp is found, it will be removed.

LENGTH OF TIME *Preparation Time ½-1 hour; Procedure ½-1 hours; Recovery ½-2 hours=2-4 Hours*

GENERAL INSTRUCTIONS

You will receive sedation for your procedure and will not be able to drive yourself home after the procedure. You must have a responsible adult accompany you home. Please have your responsible adult available to discuss your discharge instructions prior to your discharge. You will not be permitted to use any form of military or public transportation after your procedure if you are unaccompanied by an adult. Bus, taxi or shuttle drivers do not fulfill the requirement of a responsible adult.

1. ***It is recommended that females of childbearing age have a urine pregnancy test done within 72 hours prior to the procedure. Menopause is defined as last period was more than 12 months ago; if your last period was more recent, a pregnancy test will be completed on the day of the procedure.***
2. You may take Tylenol (acetaminophen), Motrin or Aspirin if you need a pain reliever. Formerly, we asked patients to stop Aspirin before their procedures. This is no longer necessary.
3. Oral iron or fiber supplements (such as Metamucil, Citrucel, Fibercon) should be discontinued 5 days prior to the exam. Please check with your physician for any substitutes.
4. Heart and blood pressure medications should be continued and taken the morning of the exam. Check with your physician regarding other medications, especially insulin, if you are diabetic.
5. If you are diabetic, hold your Metformin (Glucophage) for 24 hrs before your procedure. Hold any other oral diabetic medications the morning of the procedure and check your blood sugar before arrival. If your blood sugar is 70 or below, please inform the staff immediately upon your arrival.
6. Please report to the 6th floor of the main hospital on the day of the procedure as instructed on the day of your pre-op evaluation.
7. Leave valuables at home or with the adult who accompanies you for the procedure.
8. Wear comfortable clothes.
9. Remove all body piercings.

10. **DO NOT DRINK ANY ALCOHOL** once you begin the prep.

There will be a waiting period prior to the procedure. The WBAMC staff strives to give every patient individualized, quality care. We apologize for any inconvenience that waiting may cause you.

PROCEDURE

- 1) An intravenous line (IV) will be placed in order to provide medication for sedation and comfort.
- 2) Oxygen will be administered, and your heart and blood pressure will be continuously monitored throughout the procedure.
- 3) While lying comfortably on your left side, the physician will insert the flexible, lubricated scope into the rectum and carefully advance it through the colon. Some air and water will be placed in the colon to assist in the exam, which may be expelled afterwards. After the exam, you will rest in the recovery room until you are able to tolerate liquids and walk unassisted.

Preparation is of the utmost importance in providing accurate Colorectal Cancer Screening. The instructions below should be adhered to optimize the outcome.

2-3 Days Before your exam

Drink plenty of fluids throughout the period 2-3 days before your procedure to avoid dehydration. This consists of a minimum of 6-8 ounce glasses of fluids per day.

Avoid seeds, nuts, oatmeal or other whole grains, beans, peas, corn, and the peels of fruits and vegetables as these may remain in the colon after completion of the prep. This is a **low residue diet**.

Day before your exam

DO NOT FOLLOW ANY OTHER INSTRUCTIONS

Time	Instruction
Early am	Add water to bottle of Colyte and Refrigerate
0830 (8:30 AM)	Clear liquid breakfast
AFTER 0830 (8:30 AM)	<p>FOLLOW A RESTRICTED DIET OF CLEAR LIQUIDS</p> <ul style="list-style-type: none">• Coffee, Tea• Clear Juice without Pulp(Apple, White Grape, Lemonade)• Broth, Bouillon• Gatorade or Similar Sports Drinks• Kool-Aid, Crystal Light• Jell-o (no added fruit or toppings)• Popsicles• Carbonated Soft Drinks (Coke, Pepsi, Sprite; Diet or Regular) <p>AVOID RED OR ORANGE LIQUIDS; AVOID DAIRY PRODUCTS AND JUICE PRODUCTS WITH PULP.</p> <p>Drink plenty of fluid throughout the day to avoid dehydration. DRINK 8 OUNCES CLEAR LIQUIDS EVERY HOUR MINIMUM.</p> <p><i>The prep will work better, you will feel better and it will be easier to start your IV, if you avoid dehydration.</i></p>
1200 (NOON)	<p>Take 4 Dulcolax Tablets (if you doctor orders them).</p> <p><i>Before starting your bowel prep, make sure you are close to the restroom.</i></p>
1600 (4PM)	<ul style="list-style-type: none">• It is okay to add crystal light to improve taste of the Colyte.• It is best cold. DO NOT pour over ice. It will melt the ice and add to the volume.• Drink an 8 ounce glass every 10-15 minutes, <u>UNTIL HALF OF THE BOTTLE IS FINISHED (approx eight (8) 8 ounce glasses.)</u>• This will take you about 2 to 2 1/2 hours to drink. Some patients may feel chilled; this is not unusual.• Bowel movements should occur within 1-3 hours after the first glass.• Place the solution back in the refrigerator.• You should resume a clear liquid diet.•
0000 (Midnight)	Stop taking anything by mouth other than bowel prep (see below)

Day of the procedure

Time	Instruction
4 Hours before Leaving Home	<ul style="list-style-type: none">• Drink second half of bottle• Drink an 8 ounce glass every 10-15 minutes UNTIL THE ENTIRE BOTTLE IS COMPLETED. THIS IS CRITICAL FOR A SPLIT DOSE PREPARATION TO BE EFFECTIVE.

OUTPATIENT DISCHARGE INSTRUCTIONS:

1. You are advised to go directly home and rest quietly for the remainder of the day.
2. Until the morning after your procedure:
 - DO NOT** Drive or operate any machinery
 - DO NOT** Consume any alcoholic beverages
 - DO NOT** Sign any legal documents or make critical decisions
 - DO NOT** Take any **un-prescribed** medications
3. You may resume your normal diet.
4. A feeling of fullness or cramping from remaining air in your bowel is normal. Mild activity, such as walking, will help expel the air. Lying on your left side or directly on your stomach will also help expel the remaining air.

IF YOU HAVE ANY QUESTIONS, CONTACT THE GASTROENTEROLOGY DEPARTMENT AT 915-742-2345. PLEASE CONTACT US IF PROBLEMS ARISE BEFORE YOUR PROCEDURE OR IF YOU NEED TO CHANGE OR CANCEL YOUR PROCEDURE.

PLEASE GIVE AT LEAST 72 HOURS NOTICE WHEN CANCELING. LESS THAN 72 HOURS NOTICE MAY DELAY CARE AND INCREASE MEDICAL COST. IF YOUR PROCEDURE IS CANCELLED LESS THAN 72 HOURS PRIOR TO YOUR PROCEDURE YOU WILL BE CONSIDERED A "NO SHOW".